

OPERATORS ON SUBSPACES OF HEREDITARILY INDECOMPOSABLE BANACH SPACES

V.Ferenczi

February 1, 2008

Abstract

We show that if X is a complex hereditarily indecomposable space, then every operator from a subspace Y of X to X is of the form $\lambda I + S$, where I is the inclusion map and S is strictly singular.

1 Introduction

By *space* (resp. *subspace*), we shall always mean infinite dimensional space (resp. subspace). A space X is *hereditarily indecomposable* if no two subspaces of X are in a direct sum. In the whole article, X stands for a hereditarily indecomposable complex Banach space. It was shown in [GM] that every operator from X to X is of the form $\lambda I + S$, where I is the identity map and S is strictly singular. We generalize this result by showing that for every subspace Y of X , every operator from Y to X is of the form $\lambda I + S$, where I is the inclusion map and S is strictly singular (this was proved in [GM] in a particular case).

In fact, it is a consequence of the following lemma that it is enough to prove this for every subspace Y with a basis with constant 2.

2 Lemma

Lemma 1 *Let Y be H.I. and S an operator from Y to some Banach space. Let $Z \subset Y$. Then S is strictly singular if and only if $S|_Z$ is strictly singular.*

Proof The direct implication is clear. Now suppose that $S|_Z$ is strictly singular while S is not. Then there is a subspace Y' of Y , and $c > 0$ such that $\forall y \in Y', \|S(y)\| \geq c\|y\|$; furthermore, given $\epsilon < c/(1 + \|S\|)$, there is a subspace Z' of Z with $\|S|_{Z'}\| \leq \epsilon$. As Y is H.I., there exist two unit vectors $y \in Y', z \in Z'$ with $\|y - z\| \leq \epsilon$. Then $c - \epsilon \leq \|S(y - z)\| \leq \|S\|\epsilon$, a contradiction.

Consequence In particular, this lemma can be applied to an operator from $Y \subset X$ to X . Now suppose that the claim of the article is true for every subspace with a basis with constant 2. Let $Y \subset X$, and $T : Y \rightarrow X$. We know that Y contains a normalized basic sequence with constant 2; let Z be the subspace associated to it. By hypothesis, there exists λ such that $T|_Z - \lambda I_Z$ is strictly singular. By the lemma, $T - \lambda I_Y$ is still strictly singular.

3 A filter on block-subspaces

A normalized basic sequence $(y_i)_{i \in \mathbb{N}}$ in X will be denoted by y . We denote by $[y]$ the closed subspace generated by y , by I_y the inclusion map from $[y]$ to X , by y^k the normalized basic sequence $(y_{i+k})_{i \in \mathbb{N}}$. For an operator T such that the restriction of T to $[y]$ is defined, $T|_y$ stands for this restriction. If $(y_i)_{i \in \mathbb{N}}$ is a basic sequence such that for every i , $y_i \neq 0$, then $|y|$ denotes the normalized basic sequence $(y_i/\|y_i\|)_{i \in \mathbb{N}}$. By $y' \subset y$, we mean that y' is a normalized block basic sequence of y . Let e be a normalized basic sequence in X with constant 2. Let $J = \{y/y \subset e\}$.

Definition 1 Given y and z in J , we say that they are *similar*, and write $y \sim z$ if $\sum \|z_i - y_i\| < +\infty$.

Properties Similarity is an equivalence relation. Furthermore, for every $y \sim z$ and every $y' \subset y$, there exists $z' \subset z$ such that $y' \sim z'$. Indeed, for $k \in \mathbb{N}$, let z'_k have the same coordinates on (z_i) as y'_k on (y_i) : the normalization $|z'|$ of $(z'_k)_{k \in \mathbb{N}}$ is similar to y' .

Definition 2 For y and z in J , we say that $y \leq z$ if $\exists z' \subset z/y \sim z'$.

It is a consequence of the properties of similarity that $\leq$ is a preordering.

It is also a filter. Indeed let y and z in J ; using the fact that X is H.I., it is possible to find $y' \subset y$ and $z' \subset z$ with $y' \sim z'$. We have that $y' \leq y$ and $y' \leq z$.

Definition 3 We define a filter on the set of block subspaces of $[e]$ by letting $Y \leq Z$ if there exists $y \leq z$ with $Y = [y]$, $Z = [z]$.

Definition 4 Let $\mathcal{U}$ be an ultrafilter on J . For $y \in J$, let B_y be the Banach space of bounded operators from $[y]$ to X , and let $\|\cdot\|_y$ be the seminorm $\lim_{k \rightarrow +\infty} \|\cdot\|_{y^k}$.

Let $\mathcal{B}$ be the quotient space of $l_\infty((B_y)_{y \in J})$ by the kernel of the seminorm $\lim_{\mathcal{U}} \|\cdot\|_y$.

Lemma 2 Let $y \in J$ and $z \in J$ be similar. The operator p_{yz} from $[y]$ to $[z]$ defined by $\forall i \in \mathbb{N}, p_{yz}(y_i) = z_i$ is bounded and $\|p_{yz} - I_y\|_y = 0$.

4 A morphism from $\mathcal{L}(Y, X)$ to $\mathcal{B}$

Definition 5 Let $T : Y \rightarrow X$ with $Y = [y]$. For $z \leq y$, let $y' \subset y$ such that $z \sim y'$, let T_z be the element $Tp_{zy'}$ of B_z . Let $\tilde{T}$ be the element of $\mathcal{B}$ associated to $(T_z)_{z \leq y}$ (the value of T_z for the other values of z has no effect on the value of $\tilde{T}$, take for example $T_z = 0$).

This definition does not depend on the choice of y' . Indeed, let $y'_1 \sim y'_2$ be two choices of y' ; let T_{iz} be associated to y'_i ; then $T_{1z} - T_{2z} = T(p_{zy'_1} - p_{zy'_2})$ so that by Lemma 2, $\|T_{1z} - T_{2z}\|_z = 0$.

It is easy to check that $z \rightarrow \|T_z\|_z$ is increasing, so that $\|\tilde{T}\|$ is a simple limit.

Let B be the set of elements of $\mathcal{B}$ of the form $\tilde{T}$.

Remark 1

If $T : Y \rightarrow X$ and $U : Y \rightarrow X$, then $\widetilde{T+U} = \tilde{T} + \tilde{U}$.

By Lemma 2, for all $y \sim z$, $p_{yz} = \tilde{I}_y$.

5 Lemmas

Lemma 3 Let $T : Y \rightarrow X$ with $Y = [y]$, and let $z \leq y$. Then $\tilde{T}_z = \tilde{T}$.

Proof Let $y' \subset y$ such that $z \sim y'$. Let $z' \subset z$. For k in $\mathbb{N}$, let $t_k = p_{zy'}(z'_k)$. It has the same coordinates on (y'_i) as z'_k on (z_i) , so $z' \sim |t|$. We have

$$\sum_k \|t_k\| - 1 \leq \sum_k \|t_k - z'_k\| \leq 4 \sum_i \|y'_i - z_i\|,$$

so $\sum_k \|t_k\| - 1$ converges.

Furthermore, defining $T_{z'}'$ as $Tp_{z'|t|}$ and T_z as $Tp_{zy'}$, we get for every k :

$$\|(T_{z'}' - (T_z)_z')(z'_k)\| = \|T(|t|_k - t_k)\| \leq \|T\| \|t_k\| - 1.$$

These two points imply that $\|T_{z'}' - (T_z)_z'\|_{z'} = 0$; so $\tilde{T}_z = \tilde{T}$.

Consequences

If $y' \subset y$ then $\widetilde{T_{y'}} = \tilde{T}$.

Let $\tilde{T}$ and $\tilde{U}$ be in B , with $T : Y \rightarrow X$ and $U : Z \rightarrow X$; let $t \leq y$ and $t \leq z$; then $\tilde{T} + \tilde{U} = \tilde{T}_t + \tilde{U}_t = \widetilde{T_t + U_t}$. This proves that B is a linear space.

From now on, Y (resp. Y' , Z , Z') stands for a block subspace $[y]$ (resp. $[y']$, $[z]$, $[z']$). Recall that $Y \leq Z$ means $y \leq z$.

Proposition 1 *Let $S : Y \rightarrow X$. Then the following properties are equivalent:*

- (a) $\widetilde{S} = \widetilde{0}$.
- (b) S is strictly singular.

Proof

(a) means $\forall \epsilon > 0, \exists z \leq y$ s.t. $\|S_z\|_z \leq \epsilon$ or equivalently, the assertion:

$$(c) \forall \epsilon > 0, \exists y' \subset y, \exists k \text{ s.t. } \|S_{y',k}\| \leq 2\epsilon.$$

This last assertion implies (b) and (b) implies (c) is shown to be true in [LT].

Lemma 4 *Let $T : Y \rightarrow X$. Let Z be a block subspace of X . There exists $Y' \subset Y$, $T' : Y' \rightarrow Z$ with $\widetilde{T} = \widetilde{T'}$.*

Proof First notice that if $S : Y \rightarrow X$ is such that $\Sigma\|S(y_i)\|$ converges, then $\widetilde{S} = \widetilde{0}$.

Now let $T : Y \rightarrow X$. If $\widetilde{T} = \widetilde{0}$ then one can choose $T' = 0$, $Y' = Y$. Suppose now $\widetilde{T} \neq \widetilde{0}$. By the first consequence of Lemma 3 and Proposition 1, passing to a subspace, we may assume that T is an isomorphism.

By projections on big enough intervals, selecting block vectors, and using repeatedly the fact that X is H.I., it is possible to build a normalized block basic sequence $y' \subset y$ and a (non-normalized) block basic sequence $(z_i)_{i \in \mathbb{N}}$ in Z such that the sequence $\Sigma\|T(y'_i) - z_i\|$ converges.

The operator T' defined on $[y']$ by $T'(y'_i) = z_i$ satisfies $\widetilde{T'} = \widetilde{T}$ by the first remark, and takes its values in Z .

6 A product on B

Definition 6 Let $\widetilde{T} \in B$ with $T : Z \rightarrow X$. Let $\widetilde{U} \in B$. Let $V \leq Z$. By Lemma 4, we can assume that $\widetilde{U}$ is associated to $U : Y \rightarrow V$. We want to define $\widetilde{T}\widetilde{U}$ as $\widetilde{T_v U}$. We need to show that $\widetilde{T_v U}$ does not depend on the choice of T and U .

First, for $i = 1, 2$, let $V_i \leq Z$, and $U_i : Y_i \rightarrow V_i$ be such that $\widetilde{U}_i = \widetilde{U}$. Let $z'_i \subset z$ with $v_i \sim z'_i$. Let $y' \leq y_1$ and $y' \leq y_2$. Then

$$\widetilde{T_{v_2} U_2} - \widetilde{T_{v_1} U_1} = (\widetilde{T_{v_2} U_2})_{y'} - (\widetilde{T_{v_1} U_1})_{y'} = T_{v_2} U_{2y'} - T_{v_1} U_{1y'}.$$

Now

$$T_{v_2} U_{2y'} - T_{v_1} U_{1y'} = T(p_{v_2 z'_2} U_{2y'} - p_{v_1 z'_1} U_{1y'})$$

$$T_{v_2} U_{2y'} - T_{v_1} U_{1y'} = T[(p_{v_2 z'_2} - I_{v_2}) U_{2y'} + (U_{2y'} - U_{1y'}) + (I_{v_1} - p_{v_1 z'_1}) U_{1y'}].$$

Using Remark 1, and the fact that the space of strictly singular operators is a two-sided ideal, we get that

$$\widetilde{T_{v_2} U_2} - \widetilde{T_{v_1} U_1} = \widetilde{0}.$$

We now prove that $\widetilde{T_v U}$ does not depend on the choice of T . Let for $i = 1, 2$ $T_i : Z_i \rightarrow X$ be such that $\widetilde{T_i} = \widetilde{T}$. Let $V \leq Z_1$ and $V \leq Z_2$ and $U : Y \rightarrow V$ be a representative for $\widetilde{U}$. Then $(\widetilde{T_2})_v U - (\widetilde{T_1})_v U = ((T_2)_v - (T_1)_v)U = \widetilde{0}$.

So we can define $\widetilde{TU}$ as $\widetilde{T_v U}$ without ambiguity.

Remark 2 If $T : Z \rightarrow X$ and $U : Y \rightarrow Z$, then $\widetilde{TU} = \widetilde{TU}$. Using appropriate representatives one can then show that B is an algebra; in particular $\widetilde{1}$ the common value associated to all I_y for $y \in J$ is neutral for the multiplication.

Proposition 2 B is a field.

Proof Let $\widetilde{T} \neq \widetilde{0}$, and let us choose a representative $T : Y \rightarrow Z$ such that Z is a block subspace. By Proposition 1 and the first consequence of Lemma 3, we may assume that $T : Y \rightarrow Z$ is an isomorphism onto. Then $\widetilde{T} \widetilde{T}^{-1} = \widetilde{T T^{-1}} = \widetilde{I_{/Z}} = \widetilde{1}$; in the same way $\widetilde{T}^{-1} \widetilde{T} = \widetilde{1}$; so $\widetilde{T}$ is invertible. As B is not a singleton ($\widetilde{1} \neq \widetilde{0}$), B is a field.

7 B is a Banach algebra.

Lemma 5 For all $\lambda \in B$,

$$\|\lambda\| = \inf\{\|T\| : y \in J, T \in \mathcal{L}(Y, X), \widetilde{T} = \lambda\}.$$

Proof If $\lambda = \widetilde{0}$, the result is clear. Now assume $\lambda \neq \widetilde{0}$. For every Y , if T belongs to $\mathcal{L}(Y, X)$ then $\|\widetilde{T}\| \leq \|T_y\|_y \leq \|T\|$, so that $\|\lambda\| \leq \inf_{\widetilde{T}=\lambda} \|T\|$.

Furthermore, let $T : Z \rightarrow X$ satisfy $\lambda = \widetilde{T}$. For every $\epsilon > 0$, there exists $z' \subset z$ so that $\|T_{z'}\|_{z'} \leq (1 + \epsilon)\|\lambda\|$. So there exists k such that $\|T_{z',k}\| \leq (1 + \epsilon)^2 \|\lambda\|$. As $\widetilde{T_{z',k}} = \widetilde{T}$, we have $\inf_{\widetilde{T}=\lambda} \|T\| \leq \|\lambda\|$.

Remark We say that a representative T for $\widetilde{T}$ is ϵ -minimal if $\|T\| \leq (1 + \epsilon)\|\widetilde{T}\|$. We have proved that if T belongs to $\mathcal{L}(Y, X)$, then for any $Z \leq Y$ and any $\epsilon > 0$, we can find an ϵ -minimal representative for $\widetilde{T}$ defined on a block subspace of Z .

Proposition 3 The norm on B is an algebra norm.

Proof Let $\widetilde{T}$ and $\widetilde{U}$ be in B . Let $\epsilon > 0$. Let $T : Y \rightarrow X$ be an ϵ -minimal representative for $\widetilde{T}$. Choose an ϵ -minimal representative for $\widetilde{U}$ from some Z into Y (it is possible by the previous remark and Lemma 4).

$$\text{Then } \|\widetilde{TU}\| = \|\widetilde{TU}\| \leq \|TU\| \leq \|T\|\|U\| \leq (1 + \epsilon)^2 \|\widetilde{T}\|\|\widetilde{U}\|.$$

As ϵ is arbitrary, the norm on B is an algebra norm.

Proposition 4 *B is a Banach space.*

Proof We show that any normally converging series converges in B .

Let $(\lambda_n)_{n \in \mathbb{N}}$ be a normally converging series. Let $T_0 : Y_0 \rightarrow X$ be a 1-minimal representative for λ_0 . Given $T_{n-1} : Y_{n-1} \rightarrow X$, by the remark at the end of Lemma 5, we can find a 1-minimal representative $T_n : Y_n \rightarrow X$ for λ_n with Y_n a block subspace of Y_{n-1} .

Let $y \in J$ such that $\forall n, y_n \in Y_n$. We define an operator H on Y by $H(y_n) = \sum_{k=0}^n T_k(y_n)$. Let us evaluate $\|(H - \sum_{k=0}^n T_k)_{/y^n}\|$.

Let $u = \sum_{i=n}^{+\infty} a_i y_i$ be a vector in $[y^n]$. Then

$$\left(H - \sum_{k=0}^n T_k\right)(u) = \sum_{i=n}^{+\infty} a_i \left(\sum_{k=0}^i T_k(y_i) - \sum_{k=0}^n T_k(y_i)\right).$$

$$\left(H - \sum_{k=0}^n T_k\right)(u) = \sum_{i=n}^{+\infty} a_i \left(\sum_{k=n+1}^i T_k(y_i)\right).$$

$$\left(H - \sum_{k=0}^n T_k\right)(u) = \sum_{k=n+1}^{+\infty} T_k \left(\sum_{i=k}^{+\infty} a_i y_i\right).$$

So

$$\|(H - \sum_{k=0}^n T_k)(u)\| \leq 3 \left(\sum_{k=n+1}^{+\infty} \|T_k\|\right) \|u\|.$$

In particular, H is a bounded operator. Furthermore,

$$\lim_{n \rightarrow +\infty} \|(H - \sum_{k=0}^n T_k)_{/y^n}\| = 0.$$

By Lemma 5, $\lim_{n \rightarrow +\infty} \|\tilde{H} - \sum_{k=0}^n \lambda_k\| = 0$. This proves that $\sum_{k=0}^{+\infty} \lambda_k$ converges in B to $\tilde{H}$.

8 Conclusion

The algebra B is a Banach algebra and also a field. By Gel'fand-Mazur theorem, we can identify it to the field of complex numbers.

Now let T belong to $\mathcal{L}(Y, X)$. Then $T - \widetilde{TI_Y} = \tilde{0}$, that is $T - \widetilde{TI_Y}$ is strictly singular.

So for any block subspace Y of $[e]$, all operators on Y are of the required form. As e is arbitrary, this is true for any subspace with a basis with constant 2, and thus for any subspace.

References

- [GM] W.T. GOWERS and B. MAUREY, *The unconditional basic sequence problem*, J.Amer.Math.Soc. **6** (1993), 851-874.
- [LT] J. LINDENSTRAUSS and L. TZAFRIRI, *Classical Banach spaces I*, Springer-Verlag, New York (1977).

Equipe d'Analyse et de Mathématiques Appliquées,
Université de Marne-la-Vallée,
2, rue de la Butte Verte,
93166 Noisy le Grand Cedex, France.
ferenczi@math.univ-mlv.fr.